

Nutritional Support and Supplies

LOB(s): ⊠ Commercial	State(s): ⊠ Idaho	⊠ Washington	☐ Other:
⊠ Medicaid	□ Oregon		

Enterprise Policy

PacificSource is committed to assessing and applying current regulatory standards, widely-used treatment guidelines, and evidenced-based clinical literature when developing clinical criteria for coverage determination. Each policy contains a list of sources (references) that serves as the summary of evidence used in the development and adoption of the criteria. The evidence was considered to ensure the criteria provide clinical benefits that promote patient safety and/or access to appropriate care. Each clinical policy is reviewed, updated as needed, and readopted, at least annually, to reflect changes in regulation, new evidence, and advancements in healthcare.

Clinical Guidelines are written when necessary to provide guidance to providers and members in order to outline and clarify coverage criteria in accordance with the terms of the Member's policy. This Clinical Guideline only applies to PacificSource Health Plans, PacificSource Community Health Plans, and PacificSource Community Solutions in Idaho, Montana, Oregon, and Washington. Because of the changing nature of medicine, this list is subject to revision and update without notice. This document is designed for informational purposes only and is not an authorization or contract. Coverage determinations are made on a case-by-case basis and subject to the terms, conditions, limitations, and exclusions of the Member's policy. Member policies differ in benefits and to the extent a conflict exists between the Clinical Guideline and the Member's policy, the Member's policy language shall control. Clinical Guidelines do not constitute medical advice nor guarantee coverage.

Background

<u>Oral / Enteral nutrition</u> is indicated to maintain optimal health status for individuals with diseases or structural defects of the GI tract that interfere with transport, digestion, or absorption of nutrients. The most optimal route of nutritional intake is swallowing by mouth. In conditions where this is not possible, a tube may be placed to facilitate transport of nutrition to the digestive/absorptive sites of the GI tract. Tube placement and types are governed by individual needs, the least invasive approach being placement of a nasogastric tube. Enteral tubes may also be placed percutaneously through an abdomen. Individual needs guide the type and placement of enteral tubes, such as a nasogastric (NG) or gastrostomy tube (G-tube).

infants may qualify to receive donor human breast milk or human-derived breast milk fortifiers when human milk is not available.

To ensure the safety of donor milk, a nationally- accredited milk bank, (i.e., one using evidence-based guidelines for screening, handling, processing, and record keeping) must be used.

Regulatory Information

The term <u>Medical Food</u>, as defined in section 5(b) of the Orphan Drug Act (21 U.S.C. 360ee (b) (3)) is "a food which is formulated to be consumed or administered enterally under the supervision of a physician and which is intended for the specific dietary management of a disease or condition for which distinctive nutritional requirements, based on recognized scientific principles, are established by

medical evaluation". Medical foods are foods specially formulated for the individual who is seriously ill or who requires the product as a major treatment modality.

Non-Medical Food

Regular grocery items, over-the-counter prescription dietary products (including nutritional supplements or nutritional replacement products providing less than 60% of daily nutritional requirements), are non-medical foods and are considered not medically necessary.

Nutritional support that is taken orally (e.g., by mouth), unless mandated by state law, is not considered a medical food.

<u>Total Parenteral Nutrition (TPN)</u> is the intravenous provision of a person's complete nutritional requirements. TPN that is required for a short length of time is usually given through peripheral veins. Long-term TPN is often given through a central venous catheter (e.g., Hickman, Broviac, PICC). TPN is usually required in patients with a disease process that causes a temporary or permanent loss of absorption through the surface of the small intestine. TPN should only be an option for patients who cannot receive adequate nutritional intake via oral or enteral nutrition.

TPN can be administered safely and effectively in the patient's home by persons with specialized training. TPN is covered under the home infusion/home health benefit.

Inborn Errors of Metabolism (IEM)

Inborn errors of metabolism (IEM) include inherited biochemical disorders in which a specific enzyme defect interferes with the normal metabolism of protein, fat, or carbohydrate. As a result of diminished or absent enzyme activity in these disorders, certain compounds may accumulate in the body to toxic levels and/or other levels may become deficient. Not all disorders identified through newborn screening or other diagnostic testing require treatment with medical foods and/or dietary supplements.

The table below identifies IEMs and other diagnoses where medical foods <u>may</u> be required as part of the primary treatment.

_				
Orga	nnic	Δc	וואו	rıac

- Isovaleric acidemia
- Glutaric acidemia type I
- 3-Hydroxy 3 -methyl glutaric aciduria
- Multiple carboxylase deficiency
- Methylmalonic acidemia
- 3-Methylcrotonyl-CoA carboxylase deficiency
- · Propionic acidemia
- β-Ketothiolase

Amino Acids

- Phenylketonuria
- Maple syrup (urine) disease
- Homocystinuria
- Citrullinemia
- Argininosuccinic aciduria
- Tyrosinemia type I

Fatty Acid Oxidation

- Medium-chain acyl- CoA dehydrogenase deficiency
- Very long-chain acyl- CoA dehydrogenase deficiency
- Long-chain L-3- hydroxyacyl-CoA dehydrogenase
- Trifunctional protein deficiency
- Carnitine uptake defect

Hematology

- Sickle cell anemia
- Hb S/β thalassemia
- Hb S/C disease

Ornithine transcarbamylase deficiency

Others

 Biotinidase
 Classical galactosemia
 Congenital hypothyroid
 Congenital adrenal hyperplasia
 Critical Congenital heart disease
 Cystic fibrosis
 Glycogen storage Disease
 Guanidinoacetate
 Infantile Krabbe (Globoid Cell Leukodystrophy)
 Mucopolysaccharidosis (type 1 and type 2)
 Severe combined immunodeficiency

Criteria

Commercial

I. Oral Nutrition

Prior authorization is required

Type II (Pompe)

Spinal muscular atrophy due to

X-Linked adrenoleukodystrophy

homozygous deletion of exon 7 in SMN1

- **A.** PacificSource considers oral nutrition medically necessary when **ALL** of the following is met:
 - **1.** The requested oral nutrition product is considered a <u>Medical Food.</u> Non-Medical Food, standard infant formulas, or blenderized foods are **NOT** covered as medical food.

Note: Definitions for Medical Food and Non-Medical Food are identified in the background section of this policy.

- 2. The requested Oral nutrition product comprises 60 percent or more of the daily nutritional intake
- 3. Member has one of the diagnoses identified in the above Inborn Error of Metabolism table.

Initial authorization approval period: Up to 6 months

Reauthorization approval period: Up to 12 months

II. Enteral Nutrition via Tube

Prior authorization is required

B. PacificSource considers enteral nutrition via tube (e.g., nasogastric, jejunostomy, peg tube) feeding to be medically necessary when **ALL** of the following criteria are met:

1. The requested enteral nutrition product must be considered a Medical Food. Non-Medical Food, standard infant formulas, or blenderized foods are NOT covered as medical food.

Note: Definitions for Medical Food and Non-Medical Food are identified in the background section of this policy.

- **2.** The requested enteral nutrition product comprises <u>60 percent</u> or more of the daily nutritional intake
- **3.** The product is ordered by a physician, nurse practitioner, or registered dietician. Ordering registered dietician must work in conjunction with supervising physician or nurse practitioner.
- 4. Nutrients cannot be ingested orally due to a medical condition which either:
 - **a.** Interference with swallowing (e.g., anatomical abnormalities, dysphagia from a neurological condition, or required gavage feeding)
 - **b.** Is associated with obstruction of the proximal GI tract (e.g., tumor of the esophagus)
- **C.** PacificSource considers Digestive Enzyme that contains lipase (e.g., RELiZORB) to be medically necessary when **ALL** of the following criteria is met:
 - 1. Member is 5 years old or older
 - 2. Diagnosed with Cystic Fibrosis-Related Pancreatic Insufficiency
- **D.** Behavioral health related eating disorders- PacificSource considers enteral nutrition via tube feeding indicated for eating disorders to be medically necessary when member is unable to maintain an ideal body weight through oral feeding despite currently participating in an intensive eating disorder treatment program.

Initial authorization approval period: Up to 6 months

Reauthorization approval period: Up to 12 months

III. Donor Human Milk and Human Milk Fortifiers

- A. PacificSource considers donor human milk and/or human milk fortifiers (e.g., Prolacta) medically necessary when ordered by a licensed heath care provider or board-certified lactation consultant when ALL of the following conditions are met:
 - 1. For an <u>inpatient</u> infant who is medically or physically unable to receive maternal human milk or participate in breast feeding; **OR** for an <u>inpatient</u> infant whose parent is medically or physically unable to produce maternal human milk in sufficient quantity or caloric density, or parent is unable to participate in breast feeding
 - 2. Donor human milk must be obtained from an accredited milk bank
 - 3. The infant meets at least **ONE** of the following criteria:
 - a. Birth weight less than 2,500 grams
 - **b.** Gestational age equal to or less than 34 weeks
 - c. Hypoglycemia
 - **d.** A high risk of development of necrotizing enterocolitis, bronchopulmonary dysplasia, or retinopathy of prematurity

- **e.** A congenital or acquired gastrointestinal condition with long-term feeding or malabsorption complications
- f. Congenital heart disease requiring surgery in the first year of life
- g. An organ or bone marrow transplant
- h. Sepsis
- i. Congenital hypotonias associated with feeding difficulty or malabsorption
- j. Renal disease requiring dialysis in the first year of life
- k. Craniofacial anomalies
- I. An immunologic deficiency
- m. Neonatal abstinence syndrome
- Any other serious congenital or acquired condition for which the use of pasteurized donor human milk and donor milk derived products
- o. Any infant still inpatient within 72 hours of birth without sufficient human milk available

IV. Total Parenteral Nutrition (TPN)

Prior authorization is required

- **A.** PacificSource considers Total Parenteral Nutrition (TPN) in the home setting medically necessary when **ALL** of the following criteria are met:
 - 1. Clear documentation of a clinically significant structural or functional gastrointestinal condition, which impairs intestinal absorption
 - 2. Nutritional status cannot be maintained by **ONE** of the following:
 - **a.** Modified nutrient composition of an oral or enteral diet (e.g., lactose-free diet) due to a clinically significant structural or functional gastrointestinal condition
 - **b.** Pharmacological trial of dietary adjustment to treat the etiology of the malabsorption (e.g., pancreatic enzymes)
 - **3.** Failure of enteral nutrition as evidenced by **ONE** of the following:
 - **a.** Patient unable to receive more than 50% of daily caloric needs from oral and/or enteral diet (e.g., tube feed)
 - **b.** Serum albumin less than 3.4 grams/dL
 - c. More than 10% weight loss over a three-month (or less) period

Approved service range may be no longer than 60 days per prior authorization, up to six month maximum. Requests for more than six months or requests for lifetime TPN require Medical Director review.

Medicaid

PacificSource Community Solutions Physical Health follows Oregon Health Plan (OHP) per Oregon Administrative Rules (OAR) 410-148-0000 to 0320 for coverage of Nutritional Support and Supplies. ORS 731.162, ORS 743A.70.

PacificSource Community Solutions Pharmacy reviews CPT codes B4150 thru B4161 B9002 and B4149

Medicare

PacificSource Medicare follows Local Coverage Determination L38953 for Parenteral Nutritional Therapy and Local Coverage Determination L38955 for Enteral Nutrition.

Experimental/Investigational/Unproven

PacificSource considers digestive enzyme cartridges that connect in-line with existing enteral feeding pump tubing sets and patient extension sets or enteral feeding tubes for any other diagnosis besides Cystic Fibrosis-Related Pancreatic Insufficiency to be experimental, investigational, or unproven.

Coding Information

The following list of codes are for informational purposes only and may not be all-inclusive. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

B4100	Food Thickener, Administered Orally, Per Ounce
B4102	Enteral Formula adult fluids and electrolytes
B4103	Enteral Formula ped fluid and electrolyte
B4105	In-line cartridge containing digestive enzyme(s) for enteral feeding, each-
B4149	Enteral Formula blenderized foods
B4150	Enteral Formula Category I
B4152	Enteral Formula Category II
B4153	Enteral Formula Category III
B4154	Enteral Formula Category IV
B4155	Enteral Formula Category V
B4157	Enteral Formula special metabolic inherit
B4158	Enteral Formula ped complete intact nutrition
B4159	Enteral Formula ped complete soy based
B4160	Enteral Formula ped caloric dense >/= 0.7kc
B4161	Enteral Formula ped hydrolyzed/amino acid
B4162	Enteral Formula ped spec metabolic inherit
B4164	Parenteral nutrition solution: carbohydrates (dextrose), 50% or less ($500 \text{ ml} = 1 \text{ unit}$), home mix
B4168	Parenteral nutrition solution; amino acid, 3.5%, (500 ml = 1 unit) - home mix
B4172	Parenteral nutrition solution; amino acid, 5.5% through 7%, (500 ml = 1 unit) - home mix
B4176	Parenteral nutrition solution; amino acid, 7% through 8.5%, (500 ml = 1 unit) - home mix

B4178	Parenteral nutrition solution: amino acid, greater than 8.5% (500 ml = 1 unit) - home mix
B4180	Parenteral nutrition solution: carbohydrates (dextrose), greater than 50% (500 ml = 1 unit), home mix
B4185	Parenteral nutrition solution, not otherwise specified, 10 g lipids
B4187	Omegaven, 10 g lipids
B4189	Parenteral nutrition solution: compounded amino acid and carbohydrates with electrolytes, trace elements, and vitamins, including preparation, any strength, 10-51 g of protein, premix
B4193	Parenteral nutrition solution: compounded amino acid and carbohydrates with electrolytes, trace elements, and vitamins, including preparation, any strength, 52-73 g of protein, premix 24
B4197	Parenteral nutrition solution; compounded amino acid and carbohydrates with electrolytes, trace elements and vitamins, including preparation, any strength, 74-100 g of protein premix
B4199	Parenteral nutrition solution; compounded amino acid and carbohydrates with electrolytes, trace elements and vitamins, including preparation, any strength, over 100 g of protein, premix
B4216	Parenteral nutrition; additives (vitamins, trace elements, Heparin, electrolytes), home mix, per day
B4220	Parenteral nutrition supply kit; premix, per day
B4222	Parenteral nutrition supply kit; home mix, per day
B4224	Parenteral nutrition administration kit, per day
B5000	Parenteral nutrition solution: compounded amino acid and carbohydrates with electrolytes, trace elements, and vitamins, including preparation, any strength, renal - Amirosyn RF, NephrAmine, RenAmine – premix
B5100	Parenteral nutrition solution compounded amino acid and carbohydrates with electrolytes, trace elements, and vitamins, including preparation, any strength, hepatic-HepatAmine-premix
B5200	Parenteral nutrition solution compounded amino acid and carbohydrates with electrolytes, trace elements, and vitamins, including preparation, any strength, stress-branch chain amino acids-FreAmine-HBC-premix A
B9998	NOC for enteral supplies
B9999	Not otherwise covered parenteral supplies
S9432	Medical foods for non-inborn errors of metabolism
S9433	Medical food nutritionally complete, administered orally, providing 100% of nutritional intake
S9434	Modified solid food supplements for inborn errors of metabolism
S9435	Medical foods for inborn errors of metabolism
T2101	Human breast milk processing, storage, and distribution only
CPT® coo	les, descriptions and materials are copyrighted by the American Medical Association (AMA).
HCPCS®	codes, descriptions and materials are copyrighted by Centers for Medicare and Medicaid Services (CMS).

Definitions

Donor Human Milk - human milk contributed to a milk bank by one or more donors.

Dysphagia - difficulty swallowing due to abnormal swallowing reflex.

Inborn Errors of Metabolism - a group of rare disorders that are caused by an inherited genetic defect and alter the body's ability to derive energy from nutrients.

Intellectual Disability - Intellectual disability (ID) is a neurodevelopmental disorder that is characterized by deficits in both intellectual functioning and adaptive functioning, whose onset is in the developmental period.

Medical food - A food which is formulated to be consumed or administered enterally under the supervision of a physician, and which is intended for the specific dietary management of a disease or condition for which distinctive nutritional requirements, based on recognized scientific principles, are established by medical evaluation.

Milk Bank – an organization that engages in procurement, processing, storage, distribution, or use of human milk contributed by donors.

Related Policies

Neonatal Levels of Care and Inpatient Management

References

Advisory Committee on Heritable Disorders in Newborns and Children https://www.hrsa.gov/sites/default/files/hrsa/advisory-committees/heritable-disorders/meetings/medical-foods-report.pdf

Baker, R., D., Baker, S., S., and Bojczuk, G., (November 15, 2023), Cystic fibrosis: Nutritional issues. UpToDate. Retrieved March 22, 2023: <a href="https://www.uptodate.com/contents/cystic-fibrosis-nutritional-issues?search=Cystic+fibrosis%3A+Nutritional+issues+&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1#H29

Baker, M. L., Halliday, V., Robinson, P., Smith, K., & Bowrey, D. J. (2017). Nutrient intake and contribution of home enteral nutrition to meeting nutritional requirements after oesophagectomy and total gastrectomy. *European journal of clinical nutrition*, *71*(9), 1121–1128. https://doi.org/10.1038/ejcn.2017.88

Boullata, J. I., Clarke, J. L., Stone, A., Skoufalos, A., & Nash, D. B. (2019). Optimizing Clinical and Cost Outcomes for Patients on Enteral Nutrition Support for Treatment of Exocrine Pancreatic Insufficiency: *Proceedings from an Expert Advisory Board Meeting. Population health management*, 22(S1), S1–S10. https://doi.org/10.1089/pop.2019.0042

Camp, K. M., Lloyd-Puryear, M. A., & Huntington, K. L. (2012). Nutritional treatment for inborn errors of metabolism: indications, regulations, and availability of medical foods and dietary supplements using phenylketonuria as an example. Molecular genetics and metabolism, 107(1-2), 3–9. https://doi.org/10.1016/j.ymgme.2012.07.005

Dipasquale, V., Ventimiglia, M., Gramaglia, S., Parma, B., Funari, C., Selicorni, A., Armano, C., Salvatore, S., & Romano, C. (2019). Health-Related Quality of Life and Home Enteral Nutrition in

Children with Neurological Impairment: Report from a Multicenter Survey. *Nutrients*, *11*(12), 2968. https://doi.org/10.3390/nu11122968

Health Resources & Dervice Administration. (n.d.). Recommended uniform screening panel. HRSA. Retrieved February 22, 2023: https://www.hrsa.gov/advisory-committees/heritable-disorders/rusp

Katkin, J.P., et. al. (Sep 03, 2020). Cystic fibrosis: Assessment and management of pancreatic insufficiency. UpToDate. https://www.uptodate.com/contents/cystic-fibrosis-assessment-and-management-of-pancreatic-insufficiency?search

Medical Foods for Inborn Errors of Metabolism: History, Current Status, and Critical Need https://publications.aap.org/pediatrics/article/145/3/e20192261/36795/Medical-Foods-for-Inborn-Errors-of-Metabolism

Sheean, P., Gonzalez, M. C., Prado, C. M., McKeever, L., Hall, A. M., & Braunschweig, C. A. (2020). American Society for Parenteral and Enteral Nutrition Clinical Guidelines: The Validity of Body Composition Assessment in Clinical Populations. *JPEN. Journal of parenteral and enteral nutrition*, *44*(1), 12–43. https://doi.org/10.1002/jpen.1669

U.S. Food and Drug Administration (FDA). (March 2023). Frequently Asked Questions About Medical Foods, Third ed. Available at: https://www.fda.gov/media/97726/download. Accessed on February 29, 2024.

U.S. Food and Drug Administration (FDA). Regulatory Information. Section 5 of Orphan Drug Act. Available at:

 $\underline{http://www.fda.gov/Food/GuidanceRegulation/GuidanceDocumentsRegulatoryInformation/MedicalFoods} s/default.htm.$

U.S. Social Security Administration (SSA). (March 2017). Program Operations Manual System (POMS). DI 24598.002. Failure to Thrive. Available at: https://secure.ssa.gov/poms.nsf/lnx/0424598002.

Appendix

Policy Number:

Effective: 10/1/2020 **Next review:** 4/1/2026

Policy type: Enterprise

Author(s):

Depts.: Health Services

Applicable regulation(s): Orphan Drug Act (21 U.S.C. 360ee (b) (3)), 42 CFR §156.111; ORS 743.188, RCW 41.05.017, ,

RCW 48.43.715, MCA 33-22-131-RCW 48.44.440

Commercial OPs: 7/2025 Government OPs: 7/2025